

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ACZONE® Gel, 7.5% safely and effectively. See full prescribing information for ACZONE® Gel, 7.5%.

**ACZONE® (dapson) Gel, 7.5%, for topical use**  
**Initial U.S. Approval: 1955**

### INDICATIONS AND USAGE

ACZONE® Gel, 7.5%, is a sulfone indicated for the topical treatment of acne vulgaris in patients 12 years of age and older (1).

### DOSAGE AND ADMINISTRATION

- Apply once daily (2).
- Apply approximately a pea-sized amount of ACZONE Gel, 7.5%, in a thin layer to the entire face. A thin layer can also be applied to other affected areas (2).
- If there is no improvement after 12 weeks, treatment with ACZONE Gel, 7.5% should be reassessed (2).
- For topical use only. Not for oral, ophthalmic, or intravaginal use (2).

### DOSAGE FORMS AND STRENGTHS

Gel, 7.5% (3).

### CONTRAINDICATIONS

None (4).

### WARNINGS AND PRECAUTIONS

- Methemoglobinemia: Cases of methemoglobinemia have been reported. Discontinue ACZONE Gel if signs of methemoglobinemia occur (5.1).
- Hemolysis: Some patients with Glucose-6-phosphate Dehydrogenase (G6PD) deficiency using topical dapsone developed laboratory changes suggestive of hemolysis (5.1)(8.6).

### ADVERSE REACTIONS

Most common (incidence  $\geq$  0.9%) adverse reactions are application site dryness and pruritus (6.1).

**To report SUSPECTED ADVERSE REACTIONS, contact Allergan at 1-800-678-1605 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

- Trimethoprim/sulfamethoxazole (TMP/SMX) increases the systemic level of dapsone and its metabolites (7.1).
- Topical benzoyl peroxide used at the same time as ACZONE Gel, 7.5% may result in temporary local yellow or orange skin discoloration (7.2).

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

Revised: 05/2018

## FULL PRESCRIBING INFORMATION: CONTENTS\*

- |  |   |
|--|---|
| <b>1 INDICATIONS AND USAGE</b>                               | <b>8 USE IN SPECIFIC POPULATIONS</b>                      |
| <b>2 DOSAGE AND ADMINISTRATION</b>                           | 8.1 Pregnancy   |
| <b>3 DOSAGE FORMS AND STRENGTHS</b>                          | 8.2 Lactation   |
| <b>4 CONTRAINDICATIONS</b>                                   | 8.4 Pediatric Use   |
| <b>5 WARNINGS AND PRECAUTIONS</b>                            | 8.5 Geriatric Use   |
| 5.1 Hematological Effects                                    | 8.6 Glucose-6-phosphate Dehydrogenase (G6PD) Deficiency   |
| 5.2 Peripheral Neuropathy                                    | <b>11 DESCRIPTION</b>                                     |
| 5.3 Skin Reactions   | <b>12 CLINICAL PHARMACOLOGY</b>                           |
| <b>6 ADVERSE REACTIONS</b>                                   | 12.1 Mechanism of Action                                  |
| 6.1 Clinical Studies Experience                              | 12.3 Pharmacokinetics                                     |
| 6.2 Experience with Oral Use of Dapsone                      | 12.4 Microbiology   |
| 6.3 Postmarketing Experience                                 | <b>13 NONCLINICAL TOXICOLOGY</b>                          |
| <b>7 DRUG INTERACTIONS</b>                                   | 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility |
| 7.1 Trimethoprim-Sulfamethoxazole                            | <b>14 CLINICAL STUDIES</b>                                |
| 7.2 Topical Benzoyl Peroxide                                 | <b>16 HOW SUPPLIED/STORAGE AND HANDLING</b>               |
| 7.3 Drug Interactions with Oral Dapsone                      | <b>17 PATIENT COUNSELING INFORMATION</b>                  |
| 7.4 Concomitant Use with Drugs that Induce Methemoglobinemia |   |

\* Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

**ACZONE**<sup>®</sup> (dapsone) Gel, 7.5%, is indicated for the topical treatment of acne vulgaris in patients 12 years of age and older.

### 2 DOSAGE AND ADMINISTRATION

For topical use only. Not for oral, ophthalmic, or intravaginal use.

After the skin is gently washed and patted dry, apply approximately a pea-sized amount of **ACZONE** Gel, 7.5%, in a thin layer to the entire face once daily. In addition, a thin layer may be applied to other affected areas once daily. Rub in **ACZONE** Gel, 7.5%, gently and completely.

If there is no improvement after 12 weeks, treatment with **ACZONE** Gel, 7.5% should be reassessed (2).

### 3 DOSAGE FORMS AND STRENGTHS

Gel, 7.5%. Each gram of **ACZONE** Gel, 7.5% contains 75 mg of dapsone in an off-white to yellow gel with suspended particles.

### 4 CONTRAINDICATIONS

None.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hematological Effects

##### Methemoglobinemia

Cases of methemoglobinemia, with resultant hospitalization, have been reported postmarketing in association with twice daily dapsone gel, 5%, treatment. Patients with glucose-6-phosphate dehydrogenase deficiency or congenital or idiopathic methemoglobinemia are more susceptible to drug-induced methemoglobinemia. Avoid use of **ACZONE** Gel, 7.5% in those patients with congenital or idiopathic methemoglobinemia.

Signs and symptoms of methemoglobinemia may be delayed some hours after exposure. Initial signs and symptoms of methemoglobinemia are characterized by a slate grey cyanosis seen in e.g., buccal mucous membranes, lips, and nail beds. Advise patients to discontinue **ACZONE** Gel, 7.5% and seek immediate medical attention in the event of cyanosis.

Dapsone can cause elevated methemoglobin levels particularly in conjunction with methemoglobin-inducing agents [*see Drug Interactions (7.4)*].

##### Hemolysis

Oral dapsone treatment has produced dose-related hemolysis and hemolytic anemia. Individuals with glucose-6-phosphate dehydrogenase (G6PD) deficiency are more prone to hemolysis with the use of certain drugs. G6PD deficiency is most prevalent in populations of African, South Asian, Middle Eastern, and Mediterranean ancestry.

In clinical trials, there was no evidence of clinically relevant hemolysis or hemolytic anemia in subjects treated with topical dapsone. Some subjects with G6PD deficiency using dapsone gel, 5 %, twice daily developed laboratory changes suggestive of hemolysis [*see Use in Specific Populations (8.6)*].

Discontinue **ACZONE** Gel, 7.5%, if signs and symptoms suggestive of hemolytic anemia occur. Avoid use of **ACZONE** Gel, 7.5% in patients who are taking oral dapsone or antimalarial medications because of the potential for hemolytic reactions. Combination of **ACZONE** Gel, 7.5%, with trimethoprim/sulfamethoxazole

(TMP/SMX) may increase the likelihood of hemolysis in patients with G6PD deficiency [see Drug Interactions (7.1)].

## 5.2 Peripheral Neuropathy

Peripheral neuropathy (motor loss and muscle weakness) has been reported with oral dapsone treatment. No events of peripheral neuropathy were observed in clinical trials with topical dapsone treatment.

## 5.3 Skin Reactions

Skin reactions (toxic epidermal necrolysis, erythema multiforme, morbilliform and scarlatiniform reactions, bullous and exfoliative dermatitis, erythema nodosum, and urticaria) have been reported with oral dapsone treatment. These types of skin reactions were not observed in clinical trials with topical dapsone treatment.

## 6 ADVERSE REACTIONS

### 6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

A total of 2161 subjects were treated with **ACZONE Gel, 7.5%**, for 12 weeks in 2 controlled clinical trials. The population ranged in age from 12 to 63 years, was 56% female, and 58% Caucasian. Adverse drug reactions that were reported in at least 0.9% of subjects treated with **ACZONE Gel, 7.5%** appear in Table 1 below.

**Table 1. Adverse Reactions Occurring in at Least 0.9% of Subjects with Acne Vulgaris in 12-week Controlled Clinical Trials**

	<b>ACZONE Gel, 7.5%</b> <b>(N=2161)</b>	<b>Vehicle</b> <b>(N=2175)</b>
Application Site Dryness	24 (1.1%)	21 (1.0%)
Application Site Pruritus	20 (0.9%)	11 (0.5%)

### 6.2 Experience with Oral Use of Dapsone

Although not observed in the clinical trials with topical dapsone, serious adverse reactions have been reported with oral use of dapsone, including agranulocytosis, hemolytic anemia, peripheral neuropathy (motor loss and muscle weakness), and skin reactions (toxic epidermal necrolysis, erythema multiforme, morbilliform and scarlatiniform reactions, bullous and exfoliative dermatitis, erythema nodosum, and urticaria).

### 6.3 Postmarketing Experience

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions have been identified during post-approval use of topical dapsone: methemoglobinemia, rash (including erythematous rash, application site rash) and swelling of face (including lip swelling, eye swelling).

## 7 DRUG INTERACTIONS

No formal drug-drug interaction studies were conducted with **ACZONE Gel, 7.5%**.

### 7.1 Trimethoprim-Sulfamethoxazole

A drug-drug interaction study evaluated the effect of the use of dapsone gel, 5% in combination with double strength (160 mg/800 mg) trimethoprim-sulfamethoxazole (TMP/SMX). During co-administration, systemic levels of TMP and SMX were essentially unchanged, however, levels of dapsone and its metabolites increased in the presence of TMP/SMX. The systemic exposure from **ACZONE Gel, 7.5%** is expected to be about 1% of that from the 100 mg oral dose, even when co-administered with TMP/SMX.

### 7.2 Topical Benzoyl Peroxide

Topical application of dapsone gel followed by benzoyl peroxide in patients with acne vulgaris may result in a temporary local yellow or orange discoloration of the skin and facial hair.

### 7.3 Drug Interactions with Oral Dapsone

Certain concomitant medications (such as rifampin, anticonvulsants, St. John's wort) may increase the formation of dapsone hydroxylamine, a metabolite of dapsone associated with hemolysis. With oral dapsone treatment, folic acid antagonists such as pyrimethamine have been noted to possibly increase the likelihood of hematologic reactions.

### 7.4 Concomitant Use with Drugs that Induce Methemoglobinemia

Concomitant use of **ACZONE Gel, 7.5%** with drugs that induce methemoglobinemia such as sulfonamides, acetaminophen, acetanilide, aniline dyes, benzocaine, chloroquine, dapsone, naphthalene, nitrates and nitrites, nitrofurantoin, nitroglycerin, nitroprusside, pamaquine, para-aminosalicylic acid, phenacetin, phenobarbital, phenytoin, primaquine, and quinine may increase the risk for developing methemoglobinemia [*see Warnings and Precautions (5.1)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no available data on **ACZONE Gel, 7.5%**, use in pregnant women to inform a drug-associated risk for adverse developmental outcomes. In animal reproduction studies, oral doses of dapsone administered to pregnant rats and rabbits during organogenesis that resulted in systemic exposures more than 400 times the systemic exposure at the maximum recommended human dose (MRHD) of **ACZONE Gel, 7.5%**, resulted in embryocidal effects. When orally administered to rats from the onset of organogenesis through the end of lactation at systemic exposures approximately 500 times the exposure at the MRHD, dapsone resulted in increased stillbirths and decreased pup weight [*see Data*].

The estimated background risks of major birth defects and miscarriage for the indicated population are unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

#### Data

##### *Animal Data*

Dapsone has been shown to have an embryocidal effect in rats and rabbits when administered orally daily to females during organogenesis at dosages of 75 mg/kg/day and 150 mg/kg/day, respectively. These dosages resulted in systemic exposures that represented approximately 1407 times [rats] and 425 times [rabbits] the systemic exposure observed in human females as a result of use of the MRHD of **ACZONE Gel, 7.5%**, based on AUC comparisons. These effects were probably secondary to maternal toxicity.

Dapsone was assessed for effects on perinatal/postnatal pup development and postnatal maternal behavior and function in a study in which dapsone was orally administered to female rats daily beginning on the seventh day of gestation and continuing until the twenty-seventh day postpartum. Maternal toxicity (decreased body weight and food consumption) and developmental effects (increase in stillborn pups and decreased pup weight) were

seen at a dapsonone dose of 30 mg/kg/day (approximately 563 times the systemic exposure that is associated with the MRHD of **ACZONE** Gel, 7.5%, based on AUC comparisons). No effects were observed on the viability, physical development, behavior, learning ability, or reproductive function of surviving pups.

## 8.2 Lactation

### Risk Summary

There is no information regarding the presence of topical dapsonone in breastmilk, the effects on the breastfed infant or the effects on milk production. Orally administered dapsonone appears in human milk and could result in hemolytic anemia and hyperbilirubinemia especially in infants with G6PD deficiency. Systemic absorption of dapsonone following topical application is minimal relative to oral dapsonone administration; however, it is known that dapsonone is present in human milk following administration of oral dapsonone.

## 8.4 Pediatric Use

Safety and efficacy was evaluated in 1066 subjects aged 12-17 years old treated with **ACZONE** Gel, 7.5% in the clinical trials. The safety profile for **ACZONE** Gel, 7.5%, was similar to the vehicle control group. Safety and effectiveness of **ACZONE** Gel, 7.5%, have not been established in pediatric patients below the age of 12 years.

## 8.5 Geriatric Use

Clinical trials of **ACZONE** Gel, 7.5% did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects.


## 8.6 Glucose-6-phosphate Dehydrogenase (G6PD) Deficiency

Individuals with glucose-6-phosphate dehydrogenase (G6PD) deficiency may be more prone to methemoglobinemia and hemolysis [*see Warnings and Precautions (5.1)*].

**ACZONE** Gel, 5% and vehicle were evaluated in a randomized, double-blind, cross-over design clinical study of 64 subjects with G6PD deficiency and acne vulgaris. Subjects were Black (88%), Asian (6%), Hispanic (2%) or of other racial origin (5%). Blood samples were taken at Baseline, Week 2, and Week 12 during both vehicle and **ACZONE** Gel, 5% treatment periods. Some of these subjects developed laboratory changes suggestive of hemolysis, but there was no evidence of clinically significant hemolytic anemia in this study [*see Warnings and Precautions (5.1)*].

## 11 DESCRIPTION

**ACZONE** (dapsonone) Gel, 7.5%, contains dapsonone, a sulfone, in an aqueous gel base for topical dermatologic use. **ACZONE** Gel, 7.5% is an off-white to yellow gel with suspended particles. Chemically, dapsonone has an empirical formula of C<sub>12</sub>H<sub>12</sub>N<sub>2</sub>O<sub>2</sub>S. It is a white or slightly yellow-white, crystalline powder that has a molecular weight of 248.30. Dapsonone's chemical name is 4-[(4-aminobenzene) sulfonyl] aniline and its structural formula is:



Each gram of **ACZONE** Gel, 7.5%, contains 75 mg of dapsonone, USP, in a gel of diethylene glycol monoethyl ether, methylparaben, acrylamide/sodium acryloyldimethyl taurate copolymer, isohexadecane, polysorbate 80, and purified water.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The mechanism of action of dapsonone gel in treating acne vulgaris is not known.

### 12.3 Pharmacokinetics

In a pharmacokinetic study, male and female subjects 16 years of age or older with acne vulgaris (N=19) received 2 grams of **ACZONE** Gel, 7.5%, topically to the face, upper chest, upper back and shoulders once daily for 28 days. Steady state for dapsonone was reached within 7 days of dosing. On Day 28, the mean dapsonone maximum plasma concentration (C<sub>max</sub>) and area under the concentration-time curve from 0 to 24 hours post dose (AUC<sub>0-24h</sub>) were 13.0 ± 6.8 ng/mL and 282 ± 146 ng·h/mL, respectively. The systemic exposure from **ACZONE** Gel, 7.5% is expected to be about 1% of that from a 100 mg oral dose.

Long-term safety studies were not conducted with **ACZONE** Gel, 7.5%, however, in a long-term clinical study of dapsonone gel, 5% treatment (twice daily), periodic blood samples were collected up to 12 months to determine systemic exposure of dapsonone and its metabolites in approximately 500 subjects. Based on the measurable dapsonone concentrations from 408 subjects (M=192, F=216), obtained at Month 3, neither gender nor race appeared to affect the pharmacokinetics of dapsonone. Similarly, dapsonone exposures were approximately the same between the age groups of 12-15 years (N=155) and those greater than or equal to 16 years (N=253). There was no evidence of increasing systemic exposure to dapsonone over the study year in these subjects.

## 12.4 Microbiology

In Vivo Activity: No microbiology or immunology studies were conducted during **ACZONE** Gel, 7.5% clinical studies.

Drug Resistance: No dapsonone resistance studies were conducted during dapsonone gel clinical studies. Because no such studies were done, there are no data available as to whether dapsonone treatment may have resulted in decreased susceptibility of *Propionibacterium acnes*, an organism associated with acne, or to other antimicrobials that may be used to treat acne. Therapeutic resistance to dapsonone has been reported for *Mycobacterium leprae*, when patients have been treated with oral dapsonone.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Dapsonone was not carcinogenic to rats when orally administered to females for 92 weeks or males for 100 weeks at dose levels up to 15 mg/kg/day (approximately 340 times the systemic exposure observed in humans as a result of use of the MRHD of **ACZONE** Gel, 7.5%, based on AUC comparisons).

No evidence of potential to induce carcinogenicity was observed in a dermal study in which dapsonone gel was topically applied to Tg.AC transgenic mice for approximately 26 weeks. Dapsonone concentrations of 3%, 5%, and 10% were evaluated; 3% material was judged to be the maximum tolerated dosage.

Dapsonone was negative in a bacterial reverse mutation assay (Ames test), and was negative in a micronucleus assay conducted in mice. Dapsonone was positive (clastogenic) in a chromosome aberration assay conducted with Chinese hamster ovary (CHO) cells.

The effects of dapsonone on fertility and general reproductive performance were assessed in male and female rats following oral dosing. Dapsonone reduced sperm motility at dosages of 3 mg/kg/day or greater (approximately 22 times the systemic exposure that is associated with the MRHD of **ACZONE** Gel, 7.5%, based on AUC comparisons) when administered daily beginning 63 days prior to mating and continuing through the mating period. The mean numbers of embryo implantations and viable embryos were significantly reduced in untreated females mated with males that had been dosed at 12 mg/kg/day or greater (approximately 187 times the systemic exposure that is associated with the MRHD of **ACZONE** Gel, 7.5%, based on AUC comparisons), presumably due to reduced numbers or effectiveness of sperm, indicating impairment of fertility. When administered to female rats at a dosage of 75 mg/kg/day (approximately 1407 times the systemic exposure that is associated with the MRHD of **ACZONE** Gel, 7.5%, based on AUC comparisons) for 15 days prior to mating and for 17 days thereafter, dapsonone reduced the mean number of implantations, increased the mean early resorption rate, and reduced the mean litter size. These effects probably were secondary to maternal toxicity.



Advise the patient to read the FDA-approved patient labeling (Patient Information).

#### Hematological Effects

- Inform patients that methemoglobinemia can occur with topical dapsone treatment. Advise patients to seek immediate medical attention if they develop cyanosis [*see Warnings and Precautions (5.1)*].
- Inform patients who have G6PD deficiency that hemolytic anemia may occur with topical dapsone treatment. Advise patients to seek medical attention if they develop signs and symptoms suggestive of hemolytic anemia [*see Warnings and Precautions (5.1)*].

#### Important Administration Instructions

- Advise patients to apply **ACZONE** Gel, 7.5%, once daily to the entire face [*see Dosage and Administration (2)*].
- **ACZONE** Gel, 7.5% is for topical use only.
- Do not apply **ACZONE** Gel, 7.5% to eyes, mouth, or mucous membranes.

Distributed by: Allergan USA, Inc.  
Madison, NJ 07940

© 2018 Allergan. All rights reserved.  
All trademarks are the property of their respective owners.  
Patented. See: [www.allergan.com/patents](http://www.allergan.com/patents)



v1.0USPI5206

-----Cut Here " -----

**Patient Information**  
**ACZONE® (AK-zōn) (dapson) Gel, 7.5%**

**Important:** For use on skin only (topical use). Do not use ACZONE Gel, 7.5% in your mouth, eyes, or vagina.

**What is ACZONE Gel, 7.5%?**

ACZONE Gel, 7.5%, is a prescription medicine used on the skin (topical) to treat acne in people 12 years and older. ACZONE Gel, 7.5%, has not been studied in children under 12 years of age.

**Before you use ACZONE Gel, 7.5%, tell your doctor about all of your medical conditions, including if you:**

- have a glucose-6-phosphate dehydrogenase deficiency (G6PD)
- have higher than normal levels of methemoglobin in your blood (methemoglobinemia)
- are pregnant or plan to become pregnant. It is not known if ACZONE Gel, 7.5% will harm your unborn baby.
- are breastfeeding or plan to breastfeed. ACZONE Gel, 7.5% can pass into your breast milk and may harm your baby. You and your doctor should decide if you will use ACZONE Gel, 7.5%, or breastfeed. You should not do both.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially, tell your doctor if you are using acne medicines that contain benzoyl peroxide. Use of benzoyl peroxide with ACZONE Gel, 7.5% at the same time may cause your skin or facial hair to temporarily turn yellow or orange at the site of application.

**How do I use ACZONE Gel, 7.5%?**

- Use ACZONE Gel, 7.5% exactly as your doctor tells you to use it.
- Apply ACZONE Gel, 7.5% one time a day.
- Gently wash and pat dry the areas of your skin where you will apply ACZONE Gel, 7.5%.
- Apply a pea-sized amount of ACZONE Gel, 7.5% in a thin layer to the entire face. A thin layer may also be applied to other affected areas as instructed by your doctor.
- Rub ACZONE Gel, 7.5% in gently and completely.
- Wash your hands after applying ACZONE Gel, 7.5%.
- If your acne does not get better after using ACZONE Gel, 7.5% for 12 weeks, talk to your doctor about continuing treatment.

**What are the possible side effects of ACZONE Gel, 7.5%?**

**ACZONE Gel, 7.5% may cause serious side effects, including:**

- **Decrease of oxygen in your blood caused by a certain type of abnormal red blood cell (methemoglobinemia).** Stop using ACZONE Gel, 7.5% and get medical help right away if your lips, nail beds, or the inside of your mouth turns grey or blue.
- **Breakdown of red blood cells (hemolytic anemia).** Some people with G6PD deficiency using ACZONE Gel, 7.5% may develop mild hemolytic anemia. Stop using ACZONE Gel, 7.5% and tell your doctor right away if you get any of the following signs and symptoms:
  - back pain
  - dark brown urine
  - shortness of breath
  - fever
  - tiredness or weakness
  - yellow or pale skin

The most common side effects of ACZONE Gel, 7.5% include dryness and itching of the skin being treated.

These are not all of the possible side effects of ACZONE Gel, 7.5%. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store ACZONE Gel, 7.5%?**

- Store ACZONE Gel, 7.5%, at room temperature 68°F to 77°F (20°C to 25°C).
- Protect ACZONE Gel, 7.5% from freezing.

**Keep ACZONE Gel, 7.5% and all medicines out of the reach of children.**

**General information about the safe and effective use of ACZONE Gel, 7.5%.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ACZONE Gel, 7.5% for a condition for which it was not prescribed. Do not give ACZONE Gel, 7.5% to other people, even if they have the same symptoms you have. It may harm them. You can ask your doctor or pharmacist for information about ACZONE Gel, 7.5% that is written for health professionals.

**What are the ingredients in ACZONE Gel, 7.5%?**

**Active ingredient:** dapson

**Inactive ingredients:** diethylene glycol monoethyl ether, methylparaben, acrylamide/sodium acryloyldimethyl taurate copolymer, isohexadecane, polysorbate 80, and purified water.

Distributed by: Allergan USA, Inc.  
Madison, NJ 07940

© 2018 Allergan. All rights reserved.  
All trademarks are the property of their respective owners.  
Patented. See: [www.allergan.com/patents](http://www.allergan.com/patents)



v1.0PPI5206

For more information, call 1-800-678-1605

This Patient Information has been approved by the U.S. Food and Drug Administration.

Issued: 05/2018